

J. Raptor Res. 31(4):384–385

© 1997 The Raptor Research Foundation, Inc.

ECOLOGY OF AMERICAN KESTRELS WINTERING ON SOCORRO ISLAND, MEXICO

HARTMUT S. WALTER

University of California Los Angeles, Box 951524, Los Angeles, CA 90095-1524 U.S.A.

KEY WORDS: *American Kestrel*; *Falco sparverius*; *territoriality*; *foraging behavior*; *winter diet*.

Although American Kestrels (*Falco sparverius*) are not known to breed on the Revillagigedo Islands in the Mexican Pacific, they have been observed there in increasing numbers in recent years (Wehtje et al. 1993). I had the opportunity to observe a wintering population on Socorro Island in December 1992. This note reports on its territorial and foraging behavior as a contribution to the limited literature on the ecology of nonbreeding American Kestrels in the Neotropics.

Socorro Island lies 460 km south of Baja California's Cabo San Lucas. The main study site on Socorro (area = 140 km², highest elevation 1040 m above sea level) consisted of a 300 × 100 m grassy field at an elevation of about 450 m that was covered with grasses and weedy forbs <0.5 m in height and surrounded by low trees. Several snags (up to 2.5 m in height) of shrubby *Psidium* trees were dispersed across the open field. Nearby, there were eroded slopes, woodland patches and low scrub.

For 5 d, from the evening of 30 November to the morning of 5 December 1992, I observed the main study site and the area immediately around it for kestrels during morning and predusk hours. I observed kestrels from a distance of 40–80 m so as not to disturb them. I distinguished individual birds by their plumage and feather condition and by recording kestrels that repeatedly used and defended perch sites in hunting areas. Kestrels could not be aged with certainty. Kestrels were also seen but not monitored in other parts of the island. I did not notice any kestrels that flew to and from the island.

I counted at least 14 different kestrels in the southeast quarter of Socorro Island. Only two of them were males. The kestrels that I monitored closely showed a high degree of site fidelity. The six kestrels regularly monitored near my campsite occupied the same area day after day. One female kestrel that had its night roost in the dense foliage of a small *Bumelia* tree about 8 m away appeared to spend the entire observation time in the open, weedy field foraging in an area approximately 3 ha in size. After a few days of observations, I could accurately predict the whereabouts of this kestrel. It left its roost each morning before sunrise (0630–0645 H) and flew to one of the low snags where it hunted. Its daily routine appeared to require a minimum of energy. It would hover occasionally, then grab a small prey item from the ground and fly up

to a snag to eat it. It would stay for a few min on one snag but would then fly to another, gradually covering the entire 3 ha area. In the afternoon this female was less active. It rarely foraged in the afternoon but typically preened or soared above its territory and roost. From 1–4 December, it roosted in dense evergreen foliage at 1753, 1752, 1735 and 1722 H. Once (2 December), it roosted in the early afternoon from 1315–1720 H when it began to rain.

I observed neighboring kestrels on all sides of this kestrel. These kestrels behaved similarly and spent very little time in flight. Mostly, they perched on elevated vegetation from dawn to dusk and <10% of their time was spent in flight and foraging. All of the kestrels appeared to use their own foraging areas and I seldom observed intraspecific agonistic interactions with the exception of two birds which occasionally soared, dived and playfully chased each other in the updraft of a precipitous slope.

I observed another kestrel on 6 December from 0900–1400 H within the residential Mexican Navy compound at the southern tip of the island. It perched on metal antennas, lampposts and treetops during multiple foraging and resting periods. This bird hunted an area about 6–8 ha in size that consisted mostly of grass-covered, park-like habitat between buildings. It behaved similarly spending little time in flight.

The kestrels were obviously territorial toward other species. They aggressively defended their perches against resident Red-tailed Hawks (*Buteo jamaicensis socorroensis*) and passing Sharp-shinned Hawks (*Accipiter striatus*).

Although there were numerous small songbirds and house mice (*Mus musculus*) within the kestrel territories and the surrounding areas, I never observed a kestrel taking a bird or mammal. On several occasions, however, they captured and ate arthropod prey. The most common items were crickets, small grasshoppers, and large locusts which were abundant all over the island. Kestrels usually spotted arthropods on the ground from perches or from a low hovering position and pounced on them. No pellets were found (possibly because of the presence of large scavenging land crabs, *Gecarcinus planatus*) but I found dismembered legs of large grasshoppers or locusts at some of the perches. The apparent insectivorous diet of this population is not unexpected since breeding populations and juveniles have also been reported to be insectivorous (Balgooyen 1976, Varland et al. 1993).

Socorro Island's tropical latitude makes it a likely mi-

gration and wintering location for American Kestrels, particularly for female and juvenile birds (del Hoyo et al. 1994). The sizes of the two winter territories I observed (3 and 6–8 ha) were small compared to those observed in the northern U.S. (Craighead & Craighead 1969, Enderson 1960, Mills 1975). In California, Cade (1955) observed similarly small winter territories (e.g., a vacant lot 100 × 130 m in size). It may be that Mediterranean-type and tropical winter habitats with their mild or warm climates offer higher densities of prey biomass for wintering kestrels than do habitats in temperate and boreal climates.

RESUMEN.—Y observe lo mínimo de 14 diferente *Falco sparverius* en la isla de Socorro en el pacífico de México en 1992. Los *Falco sparverius* parecieron ocupar y defender territorios de presa en áreas herbosas en la isla donde primeramente cazaban grillos, saltamontes y langostas. Estos territorios de cazar varían en tamaño de 3–8 ha y estaban notablemente más pequeños que los antes descubiertos para *Falco sparverius*.

[Traducción de Raúl De La Garza, Jr.]

ACKNOWLEDGMENTS

Funding for the stay on Socorro Island was provided by UCLA's Latin American Center, a UC Mexus research grant and by CIBNOR in La Paz through a grant from the World Wildlife Fund (USA) for the binational Socorro Island Restoration Project (SIRP). I thank the Mexican Navy for logistic support and my research colleagues, particularly Ricardo Rodríguez-Estrella, for their assis-

tance and companionship during an unusually wet period on Socorro.

LITERATURE CITED

- BALGOOYEN, T.G. 1976. Behavior and ecology of the American Kestrel (*Falco sparverius* L.) in the Sierra Nevada of California. *Univ. Calif. Publ. Zool.* 103:1–87.
- CADE, T.J. 1955. Experiments on winter territoriality of the American Kestrel, *Falco sparverius*. *Wilson Bull.* 67: 5–17.
- CRAIGHEAD, J.J. AND F.C. CRAIGHEAD, JR. 1969. Hawks, owls and wildlife. Dover Publishers, New York, NY U.S.A.
- DEL HOYO, J., A. ELLIOTT AND J. SARGATAL [EDS.]. 1994. Handbook of the birds of the world. Vol. 2. Lynx Edicions, Barcelona, Spain.
- ENDERSON, J.H. 1960. A population study of the sparrow hawk in east-central Illinois. *Wilson Bull.* 72:222–231.
- MILLS, G.S. 1975. A winter population study of the American Kestrel in central Ohio. *Wilson Bull.* 87:241–247.
- VARLAND, D.E., E.E. KLAAS AND T.M. LOUGHIN. 1993. Use of habitat perches, causes of mortality and time until dispersal in post-fledging American Kestrels. *J. Field Ornithol.* 64:169–178.
- WEHTJE, W., H.S. WALTER, R. RODRIGUEZ-ESTRELLA, J. LLINAS AND A. CASTELLANOS-VERA. 1993. An annotated checklist of the birds of Isla Socorro, Mexico. *West. Birds* 24:1–16.

Received 12 December 1996; accepted 14 August 1997

J. Raptor Res. 31 (4):385–387

© 1997 The Raptor Research Foundation, Inc.

DIET OF THE SPECTACLED OWL (*PULSATRIX PERSPICILLATA*) DURING THE RAINY SEASON IN NORTHERN OAXACA, MEXICO

HÉCTOR GÓMEZ DE SILVA

Instituto de Ecología, UNAM, Apartado Postal 70-275, Ciudad Universitaria, UNAM, C.P. 04510, México, D.F., Mexico

MÓNICA PÉREZ-VILLAFANA

Calle 1537-3, Col. San Juan de Aragón, Sección 6, C.P. 07918, México, D.F., Mexico

JOSÉ ANTONIO SANTOS-MORENO

Departamento de Zoología, Instituto de Biología, Apartado Postal 70-153, UNAM, C.P. 04510 México, D.F., Mexico

KEY WORDS: *Spectacled Owl*; *Pulsatrix perspicillata*; *naked-tailed climbing-rat*; *Tylomys nudicaudus*; *diet*; *Oaxaca, Mexico*; *niche segregation*.

The Spectacled Owl (*Pulsatrix perspicillata*) is the largest owl in humid tropical forests of the New World, averaging

750 g in mass (Stiles and Skutch 1989). Based on its size, it is likely that it preys on the largest potential prey species in tropical forests (Emerson et al. 1994) and it is known to take mammals up to the size of agoutis (*Dasyprocta* spp.), skunks (Mephitinae) and opossums (Didelphidae). It also preys on birds as large as oropendolas (*Psar-*